

Recollements of extension algebras

CHEN Qinghua (陈清华)^{1,2} & LIN Yanan (林亚南)¹

1. Department of Mathematics, Xiamen University, Xiamen 361005, China;

2. Department of Mathematics, Fujian Normal University, Fuzhou 350007, China

Correspondence should be addressed to Lin Yanan (email: ynlin@jingxian.xmu.edu.cn)

Received January 6, 2003

Abstract Let A be a finite-dimensional algebra over arbitrary base field k . We prove: if the unbounded derived module category $D^-(\text{Mod-}A)$ admits symmetric recollement relative to unbounded derived module categories of two finite-dimensional k -algebras B and C :

$$D^-(\text{Mod} - B) \xleftarrow{\simeq} D^-(\text{Mod} - A) \xrightarrow{\simeq} D^-(\text{Mod} - C),$$

then the unbounded derived module category $D^-(\text{Mod} - T(A))$ admits symmetric recollement relative to the unbounded derived module categories of $T(B)$ and $T(C)$:

$$D^-(\text{Mod} - T(B)) \xleftarrow{\simeq} D^-(\text{Mod} - T(A)) \xrightarrow{\simeq} D^-(\text{Mod} - T(C)).$$

Keywords: trivial extension algebras, derived categories, (symmetric) recollements, partial tilting complexes.

1 Main result

As we all know, the tilting theory and the theory of derived categories are two important research subjects in the representation theory of finite-dimensional algebras, and many advanced results have been obtained in the last decades (see, for example, refs. [1–4]).

Originally, recollement for derived categories was introduced by Grothendieck in order to describe the relation between the sheaves on a topological ringed space X and those sheaves on X induced from a closed subspace and its open complement respectively^[5]. Cline et al. used the definition of recollement of triangulated categories given by Beilinson et al. in their work on perverse sheaves as a generalization of ref. [6] to obtain what they call stratification of derived module categories of certain algebras (for example, quasi-hereditary algebras)^[7]. Many mathematicians^[7–10] studied recollements of triangulated categories.

Definition^[6]. Let $\mathcal{D}, \mathcal{D}'$ and \mathcal{D}'' be triangulated categories. Then a recollement of \mathcal{D} relative to \mathcal{D}' and \mathcal{D}'' , diagrammatically expressed by

$$\mathcal{D}' \xleftarrow{\simeq} \mathcal{D} \xrightarrow{\simeq} \mathcal{D}'',$$

is given by six exact functors

$$i_* = i_! : \mathcal{D}' \rightarrow \mathcal{D}; \quad j^* = j^! : \mathcal{D} \rightarrow \mathcal{D}''; \quad i^*, i^! : \mathcal{D} \rightarrow \mathcal{D}'; \quad j_*, j_! : \mathcal{D}'' \rightarrow \mathcal{D},$$

which satisfy the following four conditions:

(R1) $(i^*, i_* = i_!, i^!)$ and $(j_!, j^* = j^!, j_*)$ are adjoint triples, i.e. i^* is left adjoint to i_* which is left adjoint to $i^!$, etc.;

(R2) $i^!j_* = 0$ (and thus $j^*i_* = 0$ and $i^*j_! = 0$);

(R3) i_* , $j_!$ and j_* are full embeddings (and thus $i^*i_* \cong i^!i_* \cong id(\mathcal{D}')$ and $j^*j_* \cong j^*j_! \cong id(\mathcal{D}'')$);

(R4) any object X in \mathcal{D} determines distinguished triangles

$$i_!i^!X \rightarrow X \rightarrow j_*j^*X \rightarrow \quad \text{and} \quad j_!j^!X \rightarrow X \rightarrow i_*i^*X \rightarrow,$$

where the morphisms $i_!i^!X \rightarrow X$ and $j_!j^!X \rightarrow X$ are the front adjunction morphisms; $X \rightarrow j_*j^*X$ and $X \rightarrow i_*i^*X$ are the rear adjunction morphisms.

Throughout this paper, all algebras are finite-dimensional algebras over arbitrary base field k . Let DA be minimal injective cogenerators of $\text{mod}A$, where $D = \text{Hom}_k(-, k)$ is the duality functor. Let $T(A)$ denote the trivial extension algebra of A . Then the trivial extension algebra $T(A)$ of A by DA is defined as the k -algebra whose additive structure is that of $A \oplus DA$ and whose multiplication structure is given by

$$(a, q)(a', q') = (aa', aq' + qa')$$

for $a, a' \in A$ and $q, q' \in DA$.

Unless stated otherwise, all our modules will be right modules. Given a k -algebra A , there are various categories of A -modules for which it will be useful to fix notation. $\text{Mod-}A$ is the category of all A -modules; $\text{mod-}A$ is the category of all finitely generated A -modules; by $D^-(\text{Mod-}A)$ we mean the derived category of unbounded complexes over $\text{Mod-}A$; by $D^b(A)$ we mean the derived category of bounded complexes over $\text{mod-}A$; $\text{Proj-}A$ is the category of all projective A -modules; P_A is the category of all finitely generated projective A -modules; by $K^b(\text{Proj-}A)$ we mean the homotopy category of bounded complexes over $\text{Proj-}A$; by $K^b(P_A)$ we mean the homotopy category of bounded complexes over P_A . Since for each algebra A there is full embedding between the various derived categories and homotopy category associated to A , we may denote homomorphisms in any of these categories just by $\text{Hom}_A(-, -)$. We shall denote by $M[n]$ rather than $T^n M$ the object obtained from a complex M by applying the "shift" functor n times. For unexplained notations, properties concerning derived categories and triangulated categories, refer to refs. [1–3], etc.

There are many important results on the recollements of triangulated categories. For convenience, we give an important Koenig's result as follows^[9].

Lemma A. Let A be a ring. Then the unbounded derived module category $D^-(\text{Mod-}A)$ admits recollement relative to the unbounded derived module categories of two rings B and C :

$$D^-(\text{Mod-}B) \xleftarrow{\quad} D^-(\text{Mod-}A) \xrightarrow{\quad} D^-(\text{Mod-}C)$$

if and only if there exist two partial-tilting complexes $M^\bullet \in K^b(\text{Proj-}A)$ and $N^\bullet \in K^b(P_A)$ that satisfy

- (1) $\text{End}_A(M^\bullet) \cong B$;
- (2) $\text{End}_A(N^\bullet) \cong C$;
- (3) $\text{Hom}_A(N^\bullet, M^\bullet) = 0$ (i.e. $\text{Hom}_A(N^\bullet, M^\bullet[n]) = 0$ for any $n \in \mathbb{Z}$);

(4) $(M^\cdot)^\perp \cap (N^\cdot)^\perp = 0$ (where $(M^\cdot)^\perp = \{Y^\cdot \in D^-(\text{Mod} - A) \mid \text{Hom}_A(M^\cdot, Y^\cdot) = 0\}$ is called the right perpendicular category determined by M^\cdot).

We say that the recollement in Lemma A is symmetric, if it satisfies the additional conditions: $\text{Hom}_A(M^\cdot, N^\cdot) = 0$ and $M^\cdot \in K^b(P_A)$. In this case, one has

$$D^-(\text{Mod} - C) \xrightleftharpoons{\sim} D^-(\text{Mod} - A) \xrightleftharpoons{\sim} D^-(\text{Mod} - B).$$

On symmetric recollement, Koenig has given some interesting characterizations^[9].

Furthermore, if $\text{gld.} A < \infty$ or $\text{gld.} C < \infty$, then $D^-(\text{Mod} - A)$ admits the recollement

$$D^-(\text{Mod} - B) \xrightleftharpoons{\sim} D^-(\text{Mod} - A) \xrightleftharpoons{\sim} D^-(\text{Mod} - C)$$

if and only if $D^b(\text{Mod} - A)$ admits recollement

$$D^b(\text{Mod} - B) \xrightleftharpoons{\sim} D^b(\text{Mod} - A) \xrightleftharpoons{\sim} D^b(\text{Mod} - C).$$

As we know, a partial tilting complex over a ring A is a complex T^\cdot in $K^b(\text{Proj} - A)$ and satisfies

- (1) $\text{Hom}_A(T^\cdot, T^\cdot[n]) = 0$ for $n \neq 0$; and
- (2) for all indexed families $\{T_i^\cdot\}_{i \in I}$ of copies of T holds:

$$\bigoplus_{i \in I} \text{Hom}_A(T^\cdot, T_i^\cdot) \xrightarrow{\text{nat.}} \text{Hom}_A(T^\cdot, \bigoplus_{i \in I} T_i^\cdot).$$

It is easy to see that partial tilting complexes are generalizations of tilting complexes and tilting complexes are generalizations of tilting modules^[2].

Our main result is the following

Theorem 1. Let A be a finite-dimensional k -algebra. If the unbounded derived module category $D^-(\text{Mod} - A)$ admits a symmetric recollement relative to the unbounded derived module categories of two finite-dimensional k -algebras B and C

$$D^-(\text{Mod} - B) \xrightleftharpoons{\sim} D^-(\text{Mod} - A) \xrightleftharpoons{\sim} D^-(\text{Mod} - C),$$

then the unbounded derived module category $D^-(\text{Mod} - T(A))$ admits symmetric recollement relative to the unbounded derived module categories of $T(B)$ and $T(C)$

$$D^-(\text{Mod} - T(B)) \xrightleftharpoons{\sim} D^-(\text{Mod} - T(A)) \xrightleftharpoons{\sim} D^-(\text{Mod} - T(C)).$$

Before we end this section, we recall the concepts of total complexes of $\text{Hom}^\cdot(X^\cdot, Y^\cdot)$ and $X^\cdot \otimes Y^\cdot$ (see ref. [11]).

Given two complexes

$$X^\cdot = \cdots \longrightarrow X^{n-1} \xrightarrow{d_X^{n-1}} X^n \xrightarrow{d_X^n} X^{n+1} \longrightarrow \cdots$$

and

$$Y^\cdot = \cdots \longrightarrow Y^{n-1} \xrightarrow{d_Y^{n-1}} Y^n \xrightarrow{d_Y^n} Y^{n+1} \longrightarrow \cdots,$$

the total complex $\text{Hom}^\cdot(X^\cdot, Y^\cdot)$ is the complex given by

$$\text{Hom}^\cdot(X^\cdot, Y^\cdot) := \cdots \longrightarrow \prod_{p \in \mathbb{Z}} \text{Hom}(X^p, Y^{l+p}) \xrightarrow{d_{\text{Hom}}^l} \prod_{p \in \mathbb{Z}} \text{Hom}(X^p, Y^{l+1+p}) \longrightarrow \cdots,$$

where $d_{\text{Hom}}^l : (\alpha^p)_{p \in \mathbb{Z}} \mapsto (d_Y^{l+p} \alpha^p - (-1)^l \alpha^{p+1} d_X^p)_{p \in \mathbb{Z}}$, for $\alpha^p \in \text{Hom}(X^p, Y^{l+p})$, and the total complex $X^\cdot \otimes Y^\cdot$ is the complex given by

$$X^\cdot \otimes Y^\cdot := \cdots \longrightarrow \bigoplus_{p \in \mathbb{Z}} X^p \otimes Y^{l-p} \xrightarrow{d_\otimes^l} \bigoplus_{p \in \mathbb{Z}} X^p \otimes Y^{l+1-p} \longrightarrow \cdots,$$

where $d_{\otimes}^l : (x \otimes y)_{p \in \mathbb{Z}} \mapsto (d_X^p(x) \otimes y + (-1)^p x \otimes d_Y^{l-p}(y))_{p \in \mathbb{Z}}$, for $x \in X^p$, $y \in Y^{l-p}$.

2 The proof of main result

Based on Lemma A, we shall give a proof of the main theorem by a series of lemmas.

Lemma 1. Let $M^\cdot = \cdots \longrightarrow M^{n-1} \xrightarrow{d_M^{n-1}} M^n \longrightarrow \cdots$ be a bounded complex of projective A -modules, i.e. $M^\cdot \in K^b(\text{Proj} - A)$. Then $R^\cdot = M^\cdot \otimes_A T(A)$ is a bounded complex of projective $T(A)$ -modules, that is, $R^\cdot \in K^b(\text{Proj} - T(A))$. Moreover, if $M^\cdot \in K^b(P_A)$, then we have $R^\cdot \in K^b(P_{T(A)})$.

Proof. For any $n \in \mathbb{Z}$, we have

$$(R^\cdot)^n = (M^\cdot \otimes_A T(A))^n = M^n \otimes_A T(A),$$

but M^\cdot is a bounded complex, so is R^\cdot .

Since any M^n is a projective A -module, supposing that $M^n = \varepsilon_n A$, where ε_n is an idempotent element of A , $M^n \otimes_{T(A)} T(A) = \varepsilon_n A \otimes_A T(A) \cong \varepsilon_n T(A)$ is a projective $T(A)$ -module.

Moreover, if any M^n is a finitely generated A -module, it is easy to know that $(R^\cdot)^n$ is also a finitely generated $T(A)$ -module.

Lemma 2. Let $M^\cdot = \cdots \longrightarrow M^{n-1} \xrightarrow{d_M^{n-1}} M^n \longrightarrow \cdots$ be a partial tilting complex of A . Then $R^\cdot = M^\cdot \otimes_A T(A)$ is a partial tilting complex of $T(A)$.

Proof. First, we prove that $\text{Hom}_{T(A)}(R^\cdot, R^\cdot[n]) = 0$ for any $n \neq 0$.

According to Iversen^[11], the homology in degree n of total complex $\text{Hom}^\cdot(R^\cdot, R^\cdot)$ is naturally isomorphic to $\text{Hom}_{T(A)}(R^\cdot, R^\cdot[n])$, so we only want to check $H^n(\text{Hom}^\cdot(R^\cdot, R^\cdot)) = 0$ for any $n \neq 0$.

According to the definition of total complex and properties of the category of complexes, and

$$\nu M^\cdot = D\text{Hom}_A(M^\cdot, A) = \text{Hom}_k(\text{Hom}_A(M^\cdot, A), k) \cong M^\cdot \otimes_A \text{Hom}_k(A, k) = M^\cdot \otimes_A DA,$$

where $\nu = D\text{Hom}(-, A)$ is the Nakayama functor, we have

$$\begin{aligned} \text{Hom}_{T(A)}^\cdot(R^\cdot, R^\cdot) &= \text{Hom}_{T(A)}^\cdot(M^\cdot \otimes_A T(A), M^\cdot \otimes_A T(A)) \\ &\cong \text{Hom}_A^\cdot(M^\cdot, \text{Hom}_{T(A)}^\cdot(T(A), M^\cdot \otimes_A T(A))) \\ &\cong \text{Hom}_A^\cdot(M^\cdot, M^\cdot \otimes_A T(A)) \\ &\cong \text{Hom}_A^\cdot(M^\cdot, M^\cdot \otimes_A (A \oplus D(A))) \\ &\cong \text{Hom}_A^\cdot(M^\cdot, M^\cdot) \oplus \text{Hom}_A^\cdot(M^\cdot, M^\cdot \otimes D(A)) \\ &\cong \text{Hom}_A^\cdot(M^\cdot, M^\cdot) \oplus \text{Hom}_A^\cdot(M^\cdot, \nu M^\cdot) \\ &\cong \text{Hom}_A^\cdot(M^\cdot, M^\cdot) \oplus D\text{Hom}_A^\cdot(M^\cdot, M^\cdot), \end{aligned}$$

hence for any $n \neq 0$,

$$H^n(\text{Hom}_{T(A)}^\cdot(R^\cdot, R^\cdot)) \cong H^n(\text{Hom}_A^\cdot(M^\cdot, M^\cdot)) \oplus H^n(D\text{Hom}_A^\cdot(M^\cdot, M^\cdot)).$$

Because M^\cdot is a partial tilting complex, for any $n \neq 0$, we have $H^n(\text{Hom}_A^\cdot(M^\cdot, M^\cdot)) = 0$, implying that $H^n(\text{Hom}_{T(A)}^\cdot(R^\cdot, R^\cdot)) = 0$. Therefore for any $n \neq 0$, $\text{Hom}_{T(A)}(R^\cdot, R^\cdot[n]) = 0$.

Second, we prove that $\oplus_{i \in I} \text{Hom}_{T(A)}(R^\cdot, R^\cdot) \stackrel{\text{nat.}}{\cong} \text{Hom}_{T(A)}(R^\cdot, \oplus_{i \in I} R_i)$, where $R_i = R^\cdot$ for

$i \in I$. In fact,

$$\begin{aligned}
 \mathrm{Hom}_{T(A)}(R, \oplus_{i \in I} R_i) &\cong \mathrm{Hom}_{T(A)}(M \otimes_A T(A), \oplus_{i \in I} (M_i \otimes_A T(A))) \\
 &\cong \mathrm{Hom}_{T(A)}(M \otimes_A T(A), (\oplus_{i \in I} M_i) \otimes_A T(A)) \\
 &\cong \mathrm{Hom}_A(M, \mathrm{Hom}_{T(A)}(T(A), (\oplus_{i \in I} M_i) \otimes_A T(A))) \\
 &\cong \mathrm{Hom}_A(M, (\oplus_{i \in I} M_i) \otimes_A T(A)) \\
 &\cong \mathrm{Hom}_A(M, \oplus_{i \in I} M_i) \oplus \mathrm{Hom}_A(M, (\oplus_{i \in I} M_i) \otimes_A DA) \\
 &\cong \mathrm{Hom}_A(M, \oplus_{i \in I} M_i) \oplus \mathrm{Hom}_A(M, \nu(\oplus_{i \in I} M_i)) \\
 &\cong \mathrm{Hom}_A(M, \oplus_{i \in I} M_i) \oplus D\mathrm{Hom}_A(\oplus_{i \in I} M_i, M) \\
 &\cong \oplus_{i \in I} \mathrm{Hom}_A(M, M_i) \oplus D \prod_{i \in I} \mathrm{Hom}_A(M_i, M)
 \end{aligned}$$

Because

$$\begin{aligned}
 D(\oplus_{i \in I} \mathrm{Hom}_A(M_i, M)) &\cong \mathrm{Hom}_k(\oplus_{i \in I} D\mathrm{Hom}_A(M_i, M), k) \\
 &\cong \prod_{i \in I} \mathrm{Hom}_k(D\mathrm{Hom}_A(M_i, M), k) \\
 &\cong \prod_{i \in I} \mathrm{Hom}_A(M_i, M),
 \end{aligned}$$

we have $\oplus_{i \in I} D\mathrm{Hom}_A(M_i, M) \cong D \prod_{i \in I} \mathrm{Hom}_A(M_i, M)$. Hence

$$\begin{aligned}
 &\mathrm{Hom}_{T(A)}(R, \oplus_{i \in I} R_i) \\
 &\cong \oplus_{i \in I} \mathrm{Hom}_A(M, M_i) \oplus (\oplus_{i \in I} D\mathrm{Hom}_A(M_i, M)) \\
 &\cong \oplus_{i \in I} (\mathrm{Hom}_A(M, M_i) \oplus D\mathrm{Hom}_A(M_i, M)) \\
 &\cong \oplus_{i \in I} (\mathrm{Hom}_A(M, M_i) \oplus \mathrm{Hom}_A(M, \nu M_i)) \\
 &\cong \oplus_{i \in I} (\mathrm{Hom}_A(M, M_i) \oplus \mathrm{Hom}_A(M, M_i \otimes_A DA)) \\
 &\cong \oplus_{i \in I} \mathrm{Hom}_A(M, M_i \otimes_A T(A)) \\
 &\cong \oplus_{i \in I} \mathrm{Hom}_{T(A)}(M \otimes_A T(A), M_i \otimes_A T(A)) \\
 &\cong \oplus_{i \in I} \mathrm{Hom}_{T(A)}(R, R_i).
 \end{aligned}$$

Lemma 3. Assume that $M \in K^b(\mathrm{Proj} - A)$ (or $M \in K^b(P_A)$), $\mathrm{End}_A(M) \cong B$. Set $R = M \otimes_A T(A)$. Then $\mathrm{End}_{T(A)}(R) \cong T(B)$.

Proof. The proof is similar to that of Theorem 3.1 in ref. [3] or Lemma 3 in ref. [12].

Lemma 4. Assume that $M \in K^b(\mathrm{Proj} - A)$, $N \in K^b(P_A)$. Set $R = M \otimes_A T(A)$, $T = N \otimes_A T(A)$. If $\mathrm{Hom}_A(N, M) = \mathrm{Hom}_A(M, N) = 0$, then $\mathrm{Hom}_{T(A)}(T, R) = \mathrm{Hom}_{T(A)}(R, T) = 0$.

Proof. Since

$$\begin{aligned}
 &\mathrm{Hom}_{T(A)}(T, R) \\
 &= \mathrm{Hom}_{T(A)}(N \otimes_A T(A), M \otimes_A T(A)) \\
 &\cong \mathrm{Hom}_A(N, \mathrm{Hom}_{T(A)}(T(A), M \otimes_A T(A))) \\
 &\cong \mathrm{Hom}_A(N, M \otimes_A T(A))
 \end{aligned}$$

$$\begin{aligned}
&\cong \text{Hom}_A(N, M) \oplus \text{Hom}_A(N, M \otimes_A D(A)) \\
&\cong \text{Hom}_A(N, \nu(M)) \quad (\text{because } \text{Hom}_A(N, M) = 0) \\
&\cong D\text{Hom}_A(M, N),
\end{aligned}$$

we have $\text{Hom}_{T(A)}(T, R) = 0$.

In the same way we can show the second equality.

Lemma 5. Let M, N, R and T be as in Lemma 4. If $(M)^\perp \cap (N)^\perp = \{0\}$, then $(R)^\perp \cap (T)^\perp = 0$.

Proof. For any $X \in (R)^\perp \cap (T)^\perp$, we have $\text{Hom}_{T(A)}(R, X[n]) = 0$ and $\text{Hom}_{T(A)}(T, X[n]) = 0$, for $n \in \mathbb{Z}$. Since

$$\begin{aligned}
\text{Hom}_{T(A)}(R, X[n]) &= \text{Hom}_{T(A)}(M \otimes_A T(A), X[n]) \\
&\cong \text{Hom}_A(M, \text{Hom}_{T(A)}(T(A), X[n])) \\
&\cong \text{Hom}_A(M, X[n]),
\end{aligned}$$

we have $\text{Hom}_A(M, X[n]) = 0$, thus $X \in (M)^\perp$.

In the same way, we have $X \in (N)^\perp$, so $X = 0$. Therefore $(R)^\perp \cap (T)^\perp = 0$.

3 Remarks

Quasi-hereditary algebras are stratification algebras; that is, their derived module category admits recollement. Indeed, quasi-hereditary algebras have the nice property that their derived module category allows a stratification with subsequent quotients isomorphic to the derived module category $D^b(k)$ of the ground field k . We point out that stratification algebras are not necessarily quasi-hereditary algebras. In order to understand the examples of algebras whose derived category admits recollement, but which are not quasi-hereditary algebras, we show first

Theorem 2. Let A, B be finite-dimensional algebras over an arbitrary base field k , ${}_A M_B$ be an A - B -bimodule, and as a k -vector space M be a finite dimension. Let $\Lambda = \begin{pmatrix} A & M \\ 0 & B \end{pmatrix}$ be the triangle matrix algebra. If $pd(M_B) < \infty$, then

$$D^b(A) \xleftarrow{\cong} D^b(\Lambda) \xleftarrow{\cong} D^b(B).$$

Proof. Let $J = \begin{pmatrix} 0 & M \\ 0 & B \end{pmatrix}$. Then J is a two-sided ideal of Λ , and $\Lambda \cong \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix} \cong A/J$ (as k -algebras). ${}_A \Lambda \cong \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} A & M \\ 0 & B \end{pmatrix} \begin{pmatrix} 1_A & 0 \\ 0 & 0 \end{pmatrix} = \Lambda \begin{pmatrix} 1_A & 0 \\ 0 & 0 \end{pmatrix}$, and $\begin{pmatrix} 1_A & 0 \\ 0 & 0 \end{pmatrix}$ is an idempotent element of Λ . Thus ${}_A \Lambda$ is a projective module. By Theorem 3.1 of ref. [8] or Theorem of ref. [10], there exists a triangulated category D'' such that $D^b(\Lambda)$ admits a recollement

$$D^b(A) \xleftarrow{\cong} D^b(\Lambda) \xleftarrow{\cong} D''.$$

In addition, $e = \begin{pmatrix} 0 & 0 \\ 0 & 1_B \end{pmatrix}$ is also an idempotent element of Λ , and $e\Lambda = \begin{pmatrix} 0 & 0 \\ 0 & B \end{pmatrix} \cong B$,

$e\Lambda e = \begin{pmatrix} 0 & 0 \\ 0 & B \end{pmatrix} \cong B$, so ${}_{e\Lambda e}e\Lambda$ is projective, hence $pd_{e\Lambda e}({}_{e\Lambda e}e\Lambda) < \infty$.

On the other hand, $\Lambda e = \begin{pmatrix} 0 & M \\ 0 & B \end{pmatrix}$; thus $\Lambda e_{e\Lambda e} \cong M_B \oplus B_B$. This implies $pd(\Lambda e_{e\Lambda e}) < \infty$.

According to Theorem 3.1 of ref. [8], there exists a triangulated category D' such that $D^b(\Lambda)$ admits a recollement

$$D' \xrightleftharpoons{\sim} D^b(\Lambda) \xrightleftharpoons{\sim} D^b(B).$$

But $\Lambda e\Lambda = \begin{pmatrix} 0 & M \\ 0 & B \end{pmatrix}$, so $\Lambda/\Lambda e\Lambda \cong A$. By Theorem of ref. [10], we have $D' \cong D^b(A)$. This completes the proof of the theorem.

Corollary 1. Let A, B be finite-dimensional algebras over an arbitrary base field k , $\Lambda = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$. Then

$$D^b(A) \xrightleftharpoons{\sim} D^b(\Lambda) \xrightleftharpoons{\sim} D^b(B).$$

Corollary 2. Let A be a finite-dimensional algebra over an arbitrary base field k , and M be a left A -module. We denote $A[M] = \begin{pmatrix} A & M \\ 0 & k \end{pmatrix}$ the one-point extension of A by M . Then

$$D^b(A) \xrightleftharpoons{\sim} D^b(A[M]) \xrightleftharpoons{\sim} D^b(k).$$

In particular, if the derived module category $D^b(A)$ of A allows successively a stratification with subsequent quotients all of the simple form $D^b(k)$, then so is the derived module category $D^b(A[M])$ of $A[M]$.

The following examples show that stratification algebras are not necessarily quasi-hereditary algebras.

Example 1. By $M_n(k)$ we denote the algebra of $n \times n$ matrices over k . Let

$$\Lambda = \left\{ \begin{pmatrix} a & 0 & 0 & e \\ 0 & b & 0 & 0 \\ 0 & d & c & 0 \\ 0 & 0 & 0 & c \end{pmatrix} \in M_4(k) \mid a, b, c, d, e \in k \right\}.$$

Then $\Lambda \cong k\vec{Q}/I$, where the quiver \vec{Q} and the relation I are given by

$$\vec{Q} = 1 \xrightarrow{\alpha} 2 \xrightarrow{\beta} 3 \quad \text{and} \quad I = \langle \alpha\beta \rangle.$$

Given the order $1 < 2 < 3$, clearly, ${}_{\Lambda}\Lambda$ is not a quasi-hereditary algebra. But if we set $A = k$,

$$B = \left\{ \begin{pmatrix} b & 0 & 0 \\ d & c & 0 \\ 0 & 0 & c \end{pmatrix} \in M_3(k) \mid b, c, d \in k \right\}, \quad M = \{(0, 0, e) \mid e \in k\}, \quad \text{then we have } \Lambda = \begin{pmatrix} A & M \\ 0 & B \end{pmatrix}$$

and $pd M_B < \infty$. So by Theorem 2, $D^b(\Lambda)$ is stratifiable with the quotients $D^b(A)$ and $D^b(B)$.

Example 2. Let A be a finite-dimensional algebra over arbitrary base field k , and $gl d A = \infty$. Set $\Lambda = \begin{pmatrix} A & A A A \\ 0 & A \end{pmatrix}$. Then by Theorem 2, we have $D^b(A) \xrightleftharpoons{\sim} D^b(\Lambda) \xrightleftharpoons{\sim} D^b(A)$; that is,

$D^b(A)$ is stratifiable with the quotients $D^b(A)$ and $D^b(A)$. But A is not a quasi-hereditary algebra for any given order.

In ref. [12] Du introduced a class of extension algebras that generalize the trivial extension algebra of A more extensively. This class of the extension algebras will be denoted by R_A^m . That is, if A is a finite-dimensional algebra over arbitrary base field k , then

$$R_A^m = \left\{ \begin{pmatrix} \lambda_1 & x_1 & 0 & \cdots & 0 & 0 \\ 0 & \lambda_2 & x_2 & \cdots & 0 & 0 \\ 0 & 0 & \lambda_3 & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & \lambda_m & x_m \\ 0 & 0 & 0 & \cdots & 0 & \lambda_1 \end{pmatrix} \mid \lambda_i \in A, x_i \in D(A), 1 \leq i \leq m \right\}.$$

Clearly, $R_A^1 = T(A)$ is a trivial extension algebra of A , and $R_A^m = \hat{A}/\langle \nu^m \rangle$, where \hat{A} is the repetitive algebra of A and ν is the Nakayama automorphism of \hat{A} . Similar to the proof of the main theorem, we have

Theorem 3. Let A be a finite-dimensional algebra, if the unbounded derived module category $D^-(\text{Mod} - A)$ admits symmetric recollement relative to the unbounded derived module categories of two finite-dimensional algebras B and C :

$$D^-(\text{Mod} - B) \xrightleftharpoons{\sim} D^-(\text{Mod} - A) \xrightleftharpoons{\sim} D^-(\text{Mod} - C),$$

then the unbounded derived category $D^-(\text{Mod} - R_A^m)$ admits symmetric recollement relative to the unbounded derived module categories of R_B^m and R_C^m :

$$D^-(\text{Mod} - R_B^m) \xrightleftharpoons{\sim} D^-(\text{Mod} - R_A^m) \xrightleftharpoons{\sim} D^-(\text{Mod} - R_C^m).$$

Acknowledgements This work was supported by the National Natural Science Foundation of China (Grant No. 10071062) and the Committee of Education Foundation of Fujian (Grant No. K2001032).

References

1. Happel, D., Triangulated categories in the representation theories of finite dimensional algebra, London Lecture Notes Series 119, New-York: Cambridge University Press, 1988.
2. Rickard, J., Morita theory for derived categories, J. London Math. Soc., 1989, 39(2): 436—456.
3. Rickard, J., Derived categories and stable equivalence, J. Pure and Appl. Algebra, 1989, 61: 303—317.
4. Hughes, D., Waschbuesch, J., Trivial extensions of tilted algebras, Proc. London Math. Soc., 1982, 46(3): 347—364.
5. Grothendieck, A., Groups and classes des categories abeliennes et trianguliers complexe parfaits, LNM589, New York: Springer-Verlag, 1977. 351—371.
6. Beilinson, A. A., Bernstein, J., Deligne, P., Faisceaux pervers, in Analyse et topologie sur les espaces singuliers, Asterisque, 1982, 100: 1—172.
7. Cline, E., Parshall, B., Scott, L., Finite dimensional algebras and hight weightest categories, J. Reine Angew Math, 1988, 391: 85—99.
8. Cline, E., Parshall, B., Scott, L., Algebraic stratification in representative categories, J. of Algebra, 1988, 117: 504—521.
9. Koenig, S., Tilting complexes, perpendicular categories and recollements of derived module categories of rings, J. Pure and Appl. Algebra, 1991, 73: 211—232.
10. Wiedemann, A., On stratificatios of derived module categories, preprint.
11. Iversen, B., Cohomology of Sheaves, Berlin: Springer-Verlag, 1986.
12. Du, X., Derived categories of algebra, Science in China, Series A, 1996, 26(11): 1002—1008.